

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

drawings are tracings made by running a pencil around the outlines of the feet and hands. The right hand is entirely normal. The left hand varies as to the little fingers, there being two digits. A careful manipulation of the palm showed that there is a single metacarpal at the base, and distally two digits, each with three bones; the bones are entirely separate, though the skin grows across between them on the level of the proximal joint. Both of the feet are six-toed, the ulnar digit in each being doubled. This doubling was found, by feeling through the skin, to begin with the digit, the metacarpal being single. The left foot bears a lump on the outside at the base of the outer digit, seemingly due to an irregularity connected with the variation presented by the toes. I was told that it had been there from before the time of fitting the first pair of shoes. There was no knowledge of a similar condition of the digits in the family; the grandparents were personally known to the young man and nothing like it had been remarked among his relatives.

The case appears to fall in the class of cases noted by Bateson ("Materials for the Study of Variation," p. 345) as among the commonest forms of polydactylism, in which there is an extra minimus, the metacarpals and metatarsals being normal. H. L. OSBORN.

5. **Pedigree Mouse Breeding.**[1] — V. Guaita has bred white and walzing mice through seven generations. There has been a loss of fecundity, due to too close inbreeding. Tables are given showing the ancestry of about three hundred mice of varying colors. The author does not make sufficient use of his data. The most striking fact is that when pure-bred white mice were crossed with pure-bred walzers (chiefly black and white), all of the twenty-eight progeny were gray, or like a house mouse, and none were walzers. When these were bred together, nine color classes at once appeared, including albinos and walzers, as well as grays and gray-blacks. If the results of breeding in the later generations are compared with what one should expect from Galton's law of ancestral inheritance, it results that the albinos appear unusually stable and prepotent. On the other hand, the walzing condition seems to be unstable and to be less potent than normal.

[1] Von Guaita, G. Zweite Mittheilung über Versuche mit Kreuzungen von verschiedenen Hausmausrassen, *Berichte d. naturf. Ges. Freiburg*, Bd. xi, Heft 2, pp. 131–138, 1899.